

---

**anti-c-Fos**

Cat #: HM1155  
Rabbit polyclonal IgG  
0.2 µg/µl, store at 4 °C

For research use only

**BACKGROUND**

The c-Fos is a cellular homolog of Fos oncogene that was detected in murine osteosarcoma virus and an avian nephroblastoma virus. It is a 62 kDa nuclear phosphoprotein that is rapidly and transiently induced by a variety of agents and functions as a transcriptional regulator for several genes. c-Fos family also include Fra-1, Fra-2 and Fos B. They are leucine zipper proteins that can dimerize with proteins of the JUN family, thereby forming the transcription factor complex AP-1. As such, the FOS proteins have been implicated as regulators of cell proliferation, differentiation, and transformation. In some cases, expression of the FOS gene has also been associated with apoptotic cell death.

**SPECIFICITY**

This antibody reacts with c-Fos, Fos B, Fra-1 and Fra-2 of mouse, rat, and human origin by Western blotting, immunoprecipitation, immunohistochemistry (including paraffin-embedded sections).

Molecular Weight of c-Fos: 55-62 kDa. Western blotting positive control: phorbol ester treated Jurkat cell lysate.

**IMMUNOGEN**

A peptide within a highly conserved domain of human c-Fos p62.

**STORAGE**

This antibody is stable for 12 months when stored at 2-8°C.

**REFERENCES**

1. Nishizawa, M., Goto, N., and Kawai, S. 1987. An avian transforming retrovirus isolated from a nephroblastoma that carries the *fos* gene as the oncogene. *J. Virol.* 61: 3733-3740.
2. Bohmann, D., Bos, T.J., Admon, A., Nishimura, T., Vogt, P.K., and Tjian, R. 1987. Human proto-oncogene *c-jun* encodes a DNA binding protein with structural and functional properties of transcription factor AP-1. *Science* 238: 1386-1392.
3. Sambucetti, L.C. and Curran, T. 1986. The *fos* protein complex is associated with DNA in isolated nuclei and binds to DNA cellulose. *Science* 234: 1417-1419.
4. Cohen, D.R., Ferreira, C.D., Gentz, R., Franza, B.R. Jr., and Curran, T. 1989. The product of a *fos*- related gene, *fra-1*, binds cooperatively to the AP-1 site with Jun: transcription factor AP-1 is comprised of multiple protein complexes. *Genes and Dev.* 3: 173-184.
5. Nishina, H., Sato, H., Suzuki, T., Sato, M., and Iba, H. 1990. Isolation and characterization of *fra-2*, an additional member of the *fos* gene family. *Proc. Natl. Acad. Sci.* 87: 3619-3623.

6. Hsu, J.C., Bravo, R. and Taub, R. (1992) Interactions among LRF-1, JunB, c-Jun, and c-Fos define a regulatory program in the G1 phase of liver regeneration. *Mol. Cell. Biol.* 12, 4654-4665.
7. Martin, M.L., Lieberman, P.M. and Curran, T. (1996) Fos-Jun dimerization promotes interaction of the basic region with TFIIE-34 and TFIIF. *Mol. Cell. Biol.* 16, 2110-2118.
8. Tanos, T., Marinissen, M.J., Leskow, F.C., Hochbaum, D., Martinetto, H., Gutkind, J.S. and Coso, O.A. (2005) Phosphorylation of c-Fos by members of the p38 MAPK family. Role in the AP-1 response to UV light. *J. Biol. Chem.* 280, 18842-18852.

**PRODUCTS FROM HYPROMATRIX, INC.****A. AntibodyArray™s:**

1. Signal Transduction AntibodyArray™  
Catalog Number HM3000
2. Apoptosis AntibodyArray™  
Catalog Number HM4000
3. Cell Cycle AntibodyArray™  
Catalog Number HM5000

**B. Staining AntibodyArray™s**

1. Staining AntibodyArray™ I  
Catalog Number HM8100
2. AntibodyArray Staining Apparatus  
Catalog Number HM8000

**C. Antibodies****1. HRP-conjugated antibodies**

- anti-phosphotyrosine  
Catalog Number HM2040
- anti-phosphoserine  
Catalog Number HM2070
- anti-phosphothreonine  
Catalog Number HM2090

and more...

**2. Primary antibodies**

Hypromatrix offers a variety of high quality antibodies. For a complete list of antibodies and their specificities, please visit our web site at [www.hypromatrix.com](http://www.hypromatrix.com).

**CONTACT**

**Hypromatrix, Inc.**  
100 Barber Avenue  
Worcester, MA 01606  
USA

Tel: 508-856-7900  
Fax: 508-302-0748  
Email: [contact@hypromatrix.com](mailto:contact@hypromatrix.com)  
Web: [www.hypromatrix.com](http://www.hypromatrix.com)